

AD-A086 207

NEW YORK UNIV N Y MAGNETO-FLUID DYNAMICS DIV F/G 12/1  
SIMILARITY SOLUTIONS OF SYSTEMS OF PARTIAL DIFFERENTIAL EQUATIO--ETC(U)  
OCT 79 P ROSENAU, J L SCHWARZMEIER AFOSR-76-2881

UNCLASSIFIED

MF-94

AFOSR-TR-80-0485

NL

[OF]  
40  
30-00000



END

DATE

FILED

8-80

DTIC

AFOSR-TR- 80-0485

COO-3077-160  
MF-94

Courant Institute of  
Mathematical Sciences

Magneto-Fluid Dynamics Division

**LEVEL II**

Similarity Solutions of Systems  
of Partial Differential  
Equations Using MACSYMA

P. Rosenau and J. L. Schwarzmeier

Air Force Office of  
Scientific Research Report  
and  
U. S. Department of Energy Report

Plasma Physics  
October 1979

New York University

Approved for public release;  
distribution unlimited.

80 7 2 078

ADA 086207

DDC FILE COPY.



DTIC  
ELECTE  
JUL 7 1980  
S C

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DDC  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 80-0485</b>	2. GOVT ACCESSION NO. <b>AD-A086207</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>SIMILARITY SOLUTIONS OF SYSTEMS OF PARTIAL DIFFERENTIAL EQUATIONS USING MACSYMA</b>	5. TYPE OF REPORT & PERIOD COVERED <b>Interim report</b>	
7. AUTHOR(s) <b>P. Rosenau J. L. Schwarzmeier</b>	8. CONTRACT OR GRANT NUMBER(s) <b>AFOSR-76-2881</b>	
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>New York University Courant Institute of Mathematical Sciences New York, NY 10012</b>	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61102F 2304/A4</b>	
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332</b>	12. REPORT DATE <b>October 1979</b>	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>MF-94, C00-3077-169</b>	13. NUMBER OF PAGES <b>24</b>	
15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  <b>Approved for public release; distribution unlimited.</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A code has been written to use the algebraic computer system MACSYMA to generate systematically the infinitesimal similarity groups corresponding to systems of quasi-linear partial differential equations. The infinitesimal similarity groups can be used to find exact solutions of the partial differential equations. In an example from fluid mechanics the similarity method using the computer code reproduces immediately a solution obtained from dimensional analysis.		

402681

JDC

UNCLASSIFIED

New York University  
Courant Institute of Mathematical Sciences  
Magneto-Fluid Dynamics Division

12

MF-94

COO-3077-160

SIMILARITY SOLUTIONS OF SYSTEMS OF  
PARTIAL DIFFERENTIAL EQUATIONS  
USING MACSYMA

P. Rosenau and J. L. Schwarzmeier


October 1979

Air Force Office of Scientific Research  
Contract No. AFOSR-76-2881

and

U. S. Department of Energy  
Contract No. EY-76-C-02-3077\*000

UNCLASSIFIED



# ABSTRACT

✓  
A code has been written to use the algebraic computer system MACSYMA to generate systematically the infinitesimal similarity groups corresponding to systems of quasi-linear partial differential equations. The infinitesimal similarity groups can be used to find exact solutions of the partial differential equations. In an example from fluid mechanics the similarity method using the computer code reproduces immediately a solution obtained from dimensional analysis.

Accession For	
NTIS	GA&I
DDC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability Codes	
Dist	Available/or special
A	

## 1. Introduction

In recent years Lie infinitesimal contact transformation group theory has become a widely used tool for systematic explorations of similarity solutions of partial differential equations [1]. Though the implementation of this method involves mostly standard manipulations (algebra and chain derivatives), in practice these become extremely involved for all but the simplest of equations.

Here we introduce the use of the algebraic computing system MACSYMA to facilitate these calculations. Specifically, MACSYMA is used to calculate systematically the generators of the infinitesimal group under which the considered equations are invariant. As far as the operational side of the similarity method is concerned, this is the most involved and tedious step. Once the general similarity group is found, a (hopefully non-trivial) subgroup is found which leaves invariant boundary curves and boundary conditions. Finally, once this subgroup is found, its invariants and consequently the similarity forms of the solutions of the partial differential equations can be found.

A primary advantage of the similarity method is that it is one of the few general techniques for obtaining exact (non-linear) solutions of partial differential equations. A primary disadvantage of the similarity method is that the solution found may satisfy only a very restricted set of initial and boundary conditions.

In general, the existence of a one parameter group of transformations leads to a reduction by one in the number of independent variables. Thus, in the most encountered case of two independent variables, the reduction leaves us with an ordinary differential equation.

We remark here that the similarity method contains as a special case those solutions which can be obtained via dimensional analysis. That is, if an equation has a solution which can be obtained from dimensional analysis, then the present method will reproduce immediately that special similarity solution, and possibly more general similarity solutions as well.

## 2. Sketch of the Similarity Method

In this section we present a brief summary of the similarity method. The similarity transformation solver (STS) uses MACSYMA to construct the similarity transformations which are admitted by a set of partial differential equations of the form (sums over  $j, k$ , and  $l$ )

$$a_{ij} \frac{\partial^2}{\partial t^2} u_j + b_{ij} \frac{\partial}{\partial t} u_j + c_{ijk} \frac{\partial}{\partial x_k} u_j + d_{ijkl} \frac{\partial^2}{\partial x_l \partial x_k} u_j = e_i, \quad (1)$$

where  $u_j = u_j(\vec{x}, t)$  are solutions, and  $a, b, c, d$ , and  $e$  are arbitrary, specified functions of  $(\vec{x}, t, \vec{u})$ . The one parameter



infinitesimal transformations considered are

$$\vec{x}' = \vec{x} + \epsilon \vec{X}(\vec{x}, t, \vec{u}) \quad (2a)$$

$$t' = t + \epsilon T(\vec{x}, t, \vec{u}) \quad (2b)$$

$$\vec{u}' = \vec{u} + \epsilon \vec{U}(\vec{x}, t, \vec{u}) \quad (2c)$$

MACSYMA is used to determine the differential equations, called the determining equations, that are satisfied by the infinitesimal groups  $\vec{X}, T, \vec{U}$ . Technically, this is achieved by requiring that the partial differential equations (1) be invariant to order  $\epsilon^2$  in terms of the new variables  $(\vec{x}', t', \vec{u}')$  of Eqs. (2). Thus the annulling of  $O(\epsilon)$  terms becomes an invariance condition.

Let  $\vec{M}[\vec{u}] = 0$  be a system of partial differential equations in the old variables, and let  $\vec{M}'[\vec{u}'] = 0$  be the same system in the new variables, where  $\vec{M}'[\vec{u}']$  is obtained from  $\vec{M}[\vec{u}]$  by replacing everywhere  $(\vec{x}, t, \vec{u}) \rightarrow (\vec{x}', t', \vec{u}')$ . The application of the transformations (2) to  $\vec{M}'[\vec{u}']$  shows that  $\vec{M}'[\vec{u}']$  is of the form  $M'[u'] = M[u] + \epsilon H + O(\epsilon^2)$ , where  $\vec{H} = \vec{H}(\vec{x}, t, \vec{u}, \vec{X}, T, \vec{U})$ . We make the system invariant to the infinitesimal transformations (2) by requiring that  $\vec{H} \equiv 0$ . Suppose we denote by  $\{S_m\}$ , for an appropriate set of values  $m$ , all possible first or second derivatives of  $u_j$  with respect to  $x_k$  and/or  $t$ . Then the expression  $\vec{H}$  is a sum of terms involving the  $S_m$ 's, with coefficients functions of  $(\vec{x}, T, \vec{U})$  and their derivatives with respect to  $(x_k, t, u_j)$ . After the original system of partial differential equations is used to

guarantee that all the  $S_m$ 's that appear in  $\vec{H}$  are independent,  $\vec{H}$  is made to vanish by requiring that all the coefficients of the  $S_m$ 's vanish. This, as explained in [1], is a sufficient but not a necessary condition. The coefficients of the  $S_m$ 's are differential equations satisfied by the infinitesimal generators  $(\vec{X}, T, \vec{U})$ ; these are the determining equations mentioned earlier. In practice the determining equations usually can be solved easily, as they are an overdetermined set of linear equations. A knowledge of the infinitesimal transformations enables one to construct a continuous transformation or in turn to find the invariants of the group that serve as the new dependent and independent variable. It is in these new variables that Eqs. (1) exhibit their symmetry and result in a reduction by one in the number of independent variables.

To illustrate the desirability of using MACSYMA to calculate the determining equations, suppose a partial differential equation contains a derivative  $\frac{\partial u}{\partial x}$  (assume scalar  $u$  and  $x$ ). In terms of the new variables, the transformed partial differential equation contains a term  $\frac{\partial u'}{\partial x}$ . But from Eq. (2c),

$$\begin{aligned} \frac{\partial u'}{\partial x'} &= \frac{\partial}{\partial x'} [u(x, t) + \epsilon U(x, t, u)] \\ &= \frac{\partial u}{\partial x} \frac{\partial x}{\partial x'} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial x'} + \epsilon \left[ \frac{\partial U}{\partial x} \frac{\partial x}{\partial x'} + \frac{\partial U}{\partial t} \frac{\partial t}{\partial x'} \right. \\ &\quad \left. + \frac{\partial U}{\partial u} \left( \frac{\partial u}{\partial x} \frac{\partial x}{\partial x'} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial x'} \right) \right] . \end{aligned} \quad (3)$$

Now for  $\frac{\partial x}{\partial x'}$  and  $\frac{\partial t}{\partial x'}$ , use Eqs. (2a-b) to obtain

$$\frac{\partial x}{\partial x'} = 1 - \epsilon \left[ \frac{\partial X}{\partial x} + \frac{\partial X}{\partial u} \frac{\partial u}{\partial x} \right] + O(\epsilon^2) \quad (4)$$

and

$$\frac{\partial t}{\partial x'} = - \epsilon \left[ \frac{\partial T}{\partial x} + \frac{\partial T}{\partial u} \frac{\partial u}{\partial x} \right] + O(\epsilon^2) \quad (5)$$

By substituting Eqs. (4) and (5) into Eq. (3) and simplifying, we obtain

$$\begin{aligned} \frac{\partial u'}{\partial x'} = \frac{\partial u}{\partial x} + \epsilon \left[ \frac{\partial u}{\partial x} + \left( \frac{\partial u}{\partial u} - \frac{\partial x}{\partial x} \right) \frac{\partial u}{\partial x} - \frac{\partial T}{\partial x} \frac{\partial u}{\partial t} - \frac{\partial x}{\partial u} \left( \frac{\partial u}{\partial x} \right)^2 \right. \\ \left. - \frac{\partial T}{\partial u} \frac{\partial u}{\partial x} \frac{\partial u}{\partial t} \right] + O(\epsilon^2) \quad (6) \end{aligned}$$

Correspondingly, any second derivatives such as  $\frac{\partial^2 u'}{\partial x'^2}$  become very involved. The system of partial differential equations (1) is written in a completely vectorized form. Since MACSYMA knows the chain rule and can simplify and factor, it is quite feasible for MACSYMA to reliably transform complicated systems of partial differential equations in several independent variables.

### 3. Programming Considerations

In this section we describe the implementation of the STS on the MACSYMA system. The basic mode of operation is that a

reference version of the STS is modified as required with the TECO editor to solve the problem at hand, and then the modified version is executed while in MACSYM using the BATCH command. Once program execution is initiated with the BATCH command, no further user interaction is required.

Input procedures for the STS are described as follows. Once a user obtains a reference copy of the STS program in his user space, a listing of the program should be made while in TECO. In the program listing the portions of the code which must be changed from one problem to another are preceded by a single asterisk and a blank line and are followed by a blank line and a double asterisk. There are three places in the STS program where the user must modify the text to solve a new problem. First, a title should be specified which identifies the system of partial differential equations being considered. Second, the user must specify the number of equations (i.e., the number of dependent variables  $u_j$ ), NEQ, the spatial dimension of the problem, NX ( $NX \geq 1$ ), and whether or not there are time derivatives ( $NT = 0$ , no time derivatives;  $NT = 1$ , there are time derivatives). Finally, the user must specify the nonzero coefficients  $a_{ij}$ ,  $b_{ij}$ ,  $c_{ijk}$ ,  $d_{ijkl}$ , and  $e_i$  or Eq. (1). Before assigning the subscript  $i$  to the coefficients, the equations should be ranked in ascending order according to the number of terms in the equations. For example, the equation with the fewest terms goes first, etc. After these changes are made in the reference STS program, the

updated version is stored with TECO as a new file, to be executed later with BATCH. In this way a user might build a library of STS programs appropriate to different systems of equations.

The output of the STS is in two stages. In the first stage the BATCH command causes listing and execution of the code. During this stage some of the intermediate calculations are printed out, so that the progress of the calculation can be inspected as it proceeds. This portion of the computing time takes about 10 minutes, nearly independent of the system of equations considered. The first intermediate printout is a listing of the functional dependencies of the functions involved in the infinitesimal transformations (2). The STS program is vectorized, so the variables  $(x_k, t, u_j, X_k, T, U_j)$  are replaced with the program variables  $(X[K], T, U[J], FX[K], FT, FU[J])$ , respectively. The second intermediate printout is a listing of the system of equations to be solved. This serves as a check that the equations were ordered properly and that the coefficients were entered correctly. Next the linearized-in-epsilon versions of the new coefficients of the transformed partial differential equations are displayed. Recall that the original system of equations must be used to guarantee that the  $S_m$ 's that appear in  $\vec{H}$  are independent. The fourth intermediate printout lists NEQ of the  $S_m$ 's which the STS has chosen to eliminate from  $\vec{H}$  by using the original system of equations. The solutions of the original equations

for the NEQ  $S_m$ 's is then listed in the last intermediate printout.

The second stage of output in the STS is the displaying of the components of  $\vec{H}$ , from which the determining equations are derived. Each component of  $\vec{H}$  is factored conveniently with respect to all the  $S_m$ 's; the determining equations can be read off immediately as the coefficients of the independent terms involving the  $S_m$ 's. The amount of time spent computing  $\vec{H}$  depends significantly on the system of equations solved, as well as on the current load on the MACSYMA system. A very rough estimate of the computing time to calculate  $\vec{H}$  is

$$T \text{ (min)} = \text{NEQ} (NX+1) (NX/2 + 2). \quad (7)$$

#### 4. An Example

As an example of the use of the STS, consider the classical Blasius problem from fluid mechanics; solve the problem of two-dimensional, steady state, incompressible, viscous, laminar flow in a boundary layer. The equations to be solved are

$$\frac{\partial}{\partial x} v_x + \frac{\partial}{\partial y} v_y = 0 \quad (8a)$$

$$v_x \frac{\partial}{\partial x} v_x + v_y \frac{\partial}{\partial y} v_x = \nu \frac{\partial^2 v_x}{\partial y^2}, \quad (8b)$$

where  $v = (v_x, v_y, 0)$  is the fluid velocity, and  $\nu$  is the coefficient of viscosity. A semi-infinite plate exists in the  $x - z$  plane for  $x \geq 0$ , and far from the plate,  $y \rightarrow \pm\infty$ ,  $v_x$

assumes a constant value,  $u_0$ . The viscous boundary conditions are:

$$v_x(y = \pm\infty) = u_0 \quad (9a)$$

$$v_x(y = 0) = v_y(y = 0) = 0. \quad (9b)$$

Using arguments of dimensional analysis, one concludes that (Landau and Lifshitz [2]) the new independent similarity variable is

$$\zeta = y \sqrt{u_0/x} \quad (10a)$$

and that  $v_x$  and  $v_y$  are of the forms

$$v_x = u_0 f_1(\zeta) \quad (10b)$$

$$v_y = \left[ \zeta/y \right] f_2(\zeta). \quad (10c)$$

Let us now obtain these same results as well as the general infinitesimal similarity group by means of the STS. In a program listing of the STS we first insert in the appropriate place a title for problem being solved. Next we insert on separate lines in the STS:

NEQ:2\$

NX:2\$

NT:0\$.

(11)

Finally we insert the nonzero coefficients:

$$\begin{aligned}
 C[1,1,1] &: 1\$ \\
 C[1,2,2] &: 1\$ \\
 C[2,1,1] &: U[1]\$ \\
 C[2,1,2] &: U[2]\$ \\
 D[2,1,2,2] &: -NU\$ ,
 \end{aligned} \tag{12}$$

where  $NU$  is the coefficient of viscosity,  $\nu$ . After this version of the STS is stored in TECO and executed in MACSYM with BATCH, the printouts begin.

The components of  $\vec{H}$  are printed out, and the determining equations can be read-off by inspection (see the Appendix). They are (letting  $U[1] \rightarrow u$ ,  $U[2] \rightarrow v$ ,  $FU[1] \rightarrow U$ , and  $FU[2] \rightarrow V$ )

$$\begin{aligned}
 X_v &= X_u = X_y = X_{vv} = Y_{vv} = Y_v \\
 &= U_{vv} = u_v = Y_u = 0 \\
 V_u - Y_x &= V_y + U_x = 0 \\
 Y_y - V_v - X_x + U_u &= U - uX_x + 2uY_y = 0 \\
 \nu (-U_{uu} + 2Y_{uy}) + 2vY_u &= -\nu U_{yy} + vU_y + uU_x = 0 \\
 \nu (Y_{yy} - 2U_{uy}) + vY_y - uY_x + V &= 0 .
 \end{aligned} \tag{13}$$

This set of differential equations for  $X, Y, U$ , and  $V$  as functions of  $x, y, u$  and  $v$  is overdetermined.



The integrations involved in solving Eqs. (13) are straightforward. The result is that the general infinitesimal similarity group corresponding to Eqs. (8) is

$$\begin{aligned} X(x,y,u,v) &= (a + 2b)x + c \\ Y(x,y,u,v) &= by + g(x) \\ U(x,y,u,v) &= au \\ V(x,y,u,v) &= ug_x - bv \end{aligned} \tag{14}$$

where  $a, b$ , and  $c$  are arbitrary constants, and  $g(x)$  is an arbitrary function of  $x$  alone.

Now we find the subgroup of (12) that leaves invariant the boundary curves (9):

$$\begin{aligned} \text{Invariance of } y = 0 &\Rightarrow y' = 0 \Rightarrow Y(x, y = 0, u, v) = 0 \\ &\Rightarrow g(x) \equiv 0. \end{aligned} \tag{15a}$$

From the transformations (2a), the boundary curves  $y = \pm \infty$  are trivially invariant.

$$\begin{aligned} \text{Invariance of } x = 0 &\Rightarrow x' = 0 \Rightarrow X(x = 0, y, u, v) = 0 \\ &\Rightarrow c = 0. \end{aligned} \tag{15b}$$

$$\begin{aligned} \text{Invariance of } x > 0 &\Rightarrow x' > 0 \Rightarrow X(x > 0, y, u, v) > 0 \\ &\Rightarrow 2b + a > 0. \end{aligned} \tag{15c}$$

Equations (15) will restrict the group (14) so that the boundary curves are left invariant. Now we further restrict

the group (14) so that the boundary conditions are invariant:

$$u(y = 0) = 0 \Rightarrow u'(y = 0) = 0 \Rightarrow U(x, y = 0, u=0, v) = 0$$

$$\Rightarrow \text{no restriction on } a . \quad (16a)$$

Similarly,  $v(y = 0) = 0$  is trivially satisfied

$$\Rightarrow \text{no restriction on } b . \quad (16b)$$

$$u(y = \pm \infty) = u_0 \Rightarrow U(x, y = \pm \infty, u = u_0, v) = 0$$

$$\Rightarrow a = 0 \quad (16c)$$

By combining Eqs. (15-16) the final subgroup of Eqs. (7-8) that leaves invariant the boundary data is

$$X(x, y, u, v) = 2bx$$

$$Y(x, y, u, v) = by \quad (17)$$

$$U(x, y, u, v) = 0$$

$$V(x, y, u, v) = -bv.$$

The first invariant surface condition is

$$\frac{dx}{X} = \frac{dy}{Y} = \frac{du}{U} , \quad \text{or,} \quad \frac{dx}{2bx} = \frac{dy}{by} = \frac{du}{0} . \quad (18)$$

Integration of the first equality in (18) is

$$\zeta = Y/\sqrt{x} = \text{constant}. \quad (19a)$$

Integration of the second equality in (18)

$$u = F_1(\zeta) . \quad (19b)$$

The second invariant surface condition is

$$\frac{dx}{x} = \frac{dy}{y} = \frac{dv}{v} , \quad \text{or,} \quad \frac{dx}{2bx} = \frac{dy}{by} = \frac{dv}{bv} . \quad (20)$$

Integration of the second equality in (20) gives

$$v = F_2(\zeta) / y . \quad (19c)$$

Equations (19) are the same results as Eqs. (10) obtained by Landau and Lifshitz. Furthermore, the infinitesimal group (14) is the general infinitesimal group corresponding to Eqs. (8). Thus the similarity solution corresponding to any other boundary conditions can be found by analyzing the group (14).

### Acknowledgements

The MACSYMA computer system is being developed by the Mathlab Group of the MIT Laboratory for Computer Science. This work is supported, in part by the Department of Energy under Contract Number E(11-1)-3070 and by the National Science Foundation under Grant NSG 1323.

### References

1. G. W. Bluman and J. D. Cole, Similarity Methods for Differential Equations, (Springer-Verlag, New York, 1974), Part II.
2. L. D. Landau and E. M. Lifshitz, Fluid Mechanics (Addison-Wesley Publishing Co., Reading, Massachusetts, 1959) pp. 145-150.

## Appendix

In this appendix we illustrate how the determining equations are obtained from a printout of the functions  $\vec{H}$ . As an example we consider the system of partial differential equations (8), so there are two components  $H_i, i = 1$  and  $2$ .

The STS factors each  $H_i$  to produce a polynomial in the  $S_m$ 's, where the factoring is performed first with respect to  $S_1$ , then with respect to  $S_2$ , and so on. That is,  $H_i$  first is factored to produce a polynomial in  $S_1$ . Then the coefficient of each power of  $S_1$  is factored to produce a polynomial in  $S_2$ , etc. In this way, once a particular product of powers of the  $S_m$ 's appears in  $H_i$ , that same combination of the  $S_m$ 's never appears again. Consequently the coefficients of the independent terms of  $H_i$  (i.e., the determining equations) can be identified by inspection.

The printout of  $\vec{H}$  corresponding to the partial differential equations (8) is

COMPONENT NUMBER 1 OF THE EXPRESSION H FROM WHICH THE  
DETERMINING EQUATIONS ARE DETERMINED IS:

$$\begin{aligned} \text{ZEROP} = & \left( \left( -\frac{d}{dU} \frac{FX}{2} - \frac{d}{dU} \frac{FX}{1} \right) * S^2 + \left( -\frac{d}{dX} \frac{FX}{2} - \frac{d}{dU} \frac{FU}{2} - \frac{d}{dX} \frac{FX}{1} + \frac{d}{dU} \frac{FU}{1} \right) * S^4 \right. \\ & + \left( \left( -\frac{d}{dU} \frac{FX}{2} - \frac{d}{dU} \frac{FX}{1} \right) * S^{13} - \frac{d}{dX} \frac{FX}{2} + \frac{d}{dU} \frac{FU}{2} \right) * S^7 \\ & \left. + \left( -\frac{d}{dX} \frac{FX}{2} + \frac{d}{dU} \frac{FU}{2} \right) * S^{13} + \frac{d}{dX} \frac{FU}{2} + \frac{d}{dX} \frac{FU}{1} \right) * \text{EPSLON} \end{aligned}$$

COMPONENT NUMBER 2 OF THE EXPRESSION H FROM WHICH THE  
DETERMINING EQUATIONS ARE DETERMINED IS:

$$\begin{aligned} \text{ZEROP} = & \left( \left( -2 * \left( -\frac{d}{dU} \frac{FX}{2} \right) * \text{NU} * S^4 + 2 * \left( -\frac{d}{dU} \frac{FX}{1} \right) * \text{NU} * S^7 + 2 * \left( -\frac{d}{dX} \frac{FX}{1} \right) * \text{NU} * S^2 \right. \right. \\ & + \left( \frac{d}{dU} \frac{FX}{2} \right) * \text{NU} * S^3 + \left( \left( -\frac{d}{dU} \frac{FX}{2} - 2 * \left( -\frac{d}{dU} \frac{FX}{1} \right) \right) * \text{NU} * S^7 \right. \\ & + \left( -2 * \left( -\frac{d}{dU} \frac{FX}{2} \right) - \frac{d}{dU} \frac{FU}{2} \right) * \text{NU} - 2 * U * \left( -\frac{d}{dU} \frac{FX}{2} \right) + U * \left( -\frac{d}{dU} \frac{FX}{1} \right) * S^2 \\ & \left. \left. + \left( \left( -2 * \left( -\frac{d}{dU} \frac{FX}{1} \right) + \frac{d}{dU} \frac{FX}{2} \right) * \text{NU} * S^2 \right. \right. \right. \end{aligned}$$

$$\begin{aligned}
& + \left( (-2 * \frac{d^2}{dU^2 dX^2} FX) + 2 * \frac{d^2}{dU dX^2} FX \right) + 2 * \frac{d^2}{dU dU^2} FU * NU - U * \frac{d}{dU^2} FX \\
& + 2 * U * \frac{d}{dU} FX * S - U * \frac{d}{dU} FX * S + \frac{d}{dU} FX * NU * S \\
& + \frac{d^2}{dX^2} FX + 2 * \frac{d^2}{dU dX^2} FU * NU + 2 * U * \frac{d}{dX^2} FX - U * \frac{d}{dX^2} FX \\
& - U * \frac{d}{dX} FX - U * \frac{d}{dU} FU + FU * S + \frac{d}{dU} FX * NU * S \\
& + \left( 2 * \frac{d^2}{dU dX^2} FX - \frac{d^2}{dU^2} FU * NU + 2 * U * \frac{d}{dU} FX * S \right) \\
& + \left( -U * \frac{d}{dU} FX * S + \frac{d}{dU} FX * NU * S + \frac{d}{dX^2} FX - 2 * \frac{d^2}{dU dX^2} FU * NU \right) \\
& + U * \frac{d}{dX^2} FX - U * \frac{d}{dX} FX + FU * S + U * \frac{d}{dU} FU * S - \frac{d}{dU} FU * NU * S \\
& - \left( \frac{d^2}{dX^2} FU * NU + U * \frac{d}{dX^2} FU + U * \frac{d}{dX} FU \right) * EPSILON
\end{aligned}$$

(D88)

BATCH DONE

(C89)



The functions  $H_i$  can be simplified considerably by making use of the following procedure: if all the independent coefficients of the  $S_m$ 's that contain only a single term are set to zero, these conditions on the groups can be used immediately to discard many of the remaining terms of  $\vec{H}$ . For instance, in  $H_2$  we note that

$$\begin{aligned} \frac{d}{dU_2} FX_1 &= \frac{d}{dU_1} FX_1 = \frac{d}{dX_2} FX_1 = \frac{d^2}{dU_2^2} FX_1 \\ &= \frac{d^2}{dU_1^2} FX_2 = \frac{d}{dU_1} FX_2 = \frac{d}{dU_2} FX_2 = \frac{d}{dU_2} FU_1 = 0 . \end{aligned} \quad (A.1)$$

By using these relationships many of the remaining terms of  $H_i$  can be crossed out immediately from the printout. For example, by using Eqs. (A.1) in  $H_1$  the terms involving  $S_4^2$ ,  $S_{13}S_7$ , and  $S_{13}$  vanish. By equating to zero the coefficients of the remaining terms of  $H_1$ , those involving  $S_4$ ,  $S_7$ , and the constant term, we obtain the following determining equations

$$\begin{aligned} \frac{d}{dX_2} FX_2 - \frac{d}{dU_2} FU_2 - \frac{d}{dX_1} FX_1 + \frac{d}{dU_1} FU_1 &= 0 , \\ - \frac{d}{dX_1} FU_2 + \frac{d}{dU_1} FU_2 &= 0 , \end{aligned}$$

and

$$\frac{d}{dx_2} FU_2 + \frac{d}{dx_1} FU_1 = 0 \quad .$$

Similar reductions occur in the size of  $H_2$  by applying  
Eqs. (A.1).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Administration, nor any person acting on behalf of the Administration:

- A. Makes any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Administration" includes any employee or contractor of the Administration, or employee of such contractor, to the extent that such employee or contractor of the Administration, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Administration, or his employment with such contractor.